

On astrophysical bounds of the cosmological constant

Andrés Balaguera-Antolínez*, Christian G. Böhrer†
and Marek Nowakowski‡

*‡Departamento de Física, Universidad de los Andes,
A.A. 4976, Bogotá, D.C., Colombia.

†Institut für Theoretische Physik, Technische Universität Wien,
Wiedner Hauptstr. 8-10, A-1040 Wien, Austria.

Abstract

Astrophysical bounds on the cosmological constant are examined for spherically symmetric bodies. Similar limits emerge from hydrostatical and gravitational equilibrium and the validity of the Newtonian limit. The methods in use seem to be disjoint from the basic principles, however they have the same implication regarding the upper bounds. Therefore we will compare different inequalities and comment on the possible relationship between them. These inequalities are of relevance for the so-called coincidence problem and for the bound of the cosmological constant which comes surprisingly close to the ‘experimental’ value.

Keywords: cosmological constant, relativistic astrophysical objects, virial theorem

PACS numbers: 95.30.Sf, 04.40.Dg

*e-mail: a-balagu@uniandes.edu.co

†e-mail: boehmer@hep.itp.tuwien.ac.at

‡e-mail: mnowakos@uniandes.edu.co

1 Introduction

The cosmological constant Λ has a history as long as General Relativity itself: introduced to make the universe static, abandoned after the discovery of the Hubble expansion and introduced recently to explain the acceleration of the Universe [1, 2]. It is true that the latter phenomenon can, in principle, be also accounted for by other explanations like quintessence [3, 4], Chaplygin Gas [5], mixture models [6–9] and modified gravity [10] etc, but the simplest explanation for an acceleration can be obtained from a positive cosmological constant with

$$\begin{aligned}\Lambda &= 8\pi G\rho_{\text{vac}}, \\ \rho_{\text{vac}} &\approx (0.7 - 0.8)\rho_{\text{crit}}.\end{aligned}\tag{1}$$

This value is surprisingly close to a prediction made long ago by Zel’dovich [11–14], namely,

$$\Lambda \approx G^2 m^6,\tag{2}$$

where m is a typical hadronic mass scale ($0.15 - 1$ GeV). Indeed, combining this with $H_0 = m^3$ [15, 16], one gets the result that the Hubble parameter is dominated by the cosmological constant Λ .

Cosmological observations [17–19] give strong indications of the presence of a positive cosmological constant which would mean that the universe is of de Sitter type. Moreover, recent studies [20] of X-Rays hint toward a constant density of the *dark energy* which is thought responsible for the acceleration, also in agreement with a positive Λ . The use of the different distance measurements in a Λ CDM model (models of cold dark matter with Λ) also shows strong evidence for a positive cosmological constant ($0.47 < \Omega_\Lambda < 0.82$, $\Omega_\Lambda = \rho_{\text{vac}}/\rho_{\text{crit}}$) [21].

Such a relatively large cosmological constant has astrophysical implications. This was realized as early as 1939 by Gamow and Teller [22]. The situation is best paraphrased in [23] which summarizes the work of several authors in the 30’s and 40’s: “The essential difficulty with a relativistic theory in which λ [the Cosmological Constant] is positive is that of accounting for the formation and condensation in terms of gravitational instability; for, to use the ‘*force*’ metaphor, the present expansion indicates that the force of cosmic repulsion exceeds those of gravitational attraction [22]. This is not likely to disturb the stability of systems (such as the galaxy) of high average density, but it is likely to prevent new condensation in regions of low density.”

Indeed, several studies have corroborated this conclusion and many well known systems are reconsidered with the inclusion of the formerly omitted Λ -term. Another, more mathematical reason to include Λ is the richer asymptotic structure of these spacetimes and moreover the AdS/CFT correspondence [24] motivating much research.

Examples of recent research are the Einstein-Yang-Mills- Λ system [25], ($n + 4$) dimensional EYM- Λ theory [26], perfect fluid solutions with cosmological constant [27–30], consistency with the Newtonian limit [31, 32], gravitational

equilibrium via the virial theorem [33] and also Einstein-Cartan and Einstein-Cartan-Dirac theory with cosmological constant [34, 35]. It is remarkable that these examples have something in common, namely, in each of them there exists an upper bound of the cosmological constant.

Gravitational equilibrium and perfect fluid considerations surprisingly predict the same upper bound in terms of the mean density of an astrophysical object, $\Lambda < 4\pi G\bar{\rho}$. Hence it seems that there is some relation between the existence of exact solutions of Einstein's field equations and the virial theorem. In the EYM- Λ system, one finds $\Lambda < \Lambda_{\text{crit}}$. In the $(n+4)$ dimensional EYM- Λ theory it was found that the cosmological constant is bounded, such that $\Lambda_{(n+4)} < g^2/2G_{(n+4)}$ and constant density solutions exist for $\Lambda < \Lambda_S$, for larger values they are unphysical. Finally we mention that supergravity [36] even forbids a positive value and therefore predicts $\Lambda \leq 0$. It should be noted that a lower bound never occurs.

In [33] it was found that the more the shape of the object deviates from spherical symmetry, the more difficult it is to reach the gravitational equilibrium in presence of a positive cosmological constant for low density astrophysical objects. This result was based on the assumption of gravitational equilibrium. Of course, equally valid and useful concept is the hydrodynamical equilibrium combined with Einstein's equations. These two concepts of equilibrium give us information on Λ in terms of inequalities. A third source of useful information on Λ is the Newtonian limit, which also gives us an inequality.

In this paper we will compare different inequalities and point out some interesting coincidences. We will use these results to infer on certain static properties of astrophysical objects. Unless otherwise stated we put $G = c = \hbar = 1$.

2 Newtonian hydrostatic equilibrium

In order to be self consistent, some basic relations are reviewed as shortly as possible.

In Newtonian astrophysics a spherically symmetric object is stable if the gravitational and matter pressure are in equilibrium. This condition leads to

$$P'(r) = -r\rho(r)\left(\frac{m(r)}{r^3} - \frac{\Lambda}{3}\right), \quad (3)$$

which is sometimes called [37, p.301] the "fundamental equation of Newtonian astrophysics". The mass function is as usual defined by

$$m(r) = \int_0^r 4\pi\rho(s)s^2 ds. \quad (4)$$

Furthermore let the mean density up to r be defined by

$$\bar{\rho}(r) = \frac{3}{4\pi} \frac{m(r)}{r^3}, \quad (5)$$

which then has the following properties

$$\bar{\rho}_c = \rho_c, \quad \bar{\rho}(r) \geq \rho(r) \quad \forall r, \quad (6)$$

where ρ_c is the central density. If a density profile $\rho(r)$ for the astrophysical object is prescribed, one can firstly use (4) to find the mass function and secondly integrate (3) to find the pressure $P(r)$.

However the most physical starting point is to prescribe an equation of state $\rho = \rho(P)$. First we rewrite (3) with the help of (5) and find

$$P'(r) = -r \frac{\rho(r)}{3} (4\pi \bar{\rho}(r) - \Lambda). \quad (7)$$

Then, differentiating the mean density (5) leads to

$$\bar{\rho}'(r) = \frac{3}{r} (\rho(P(r)) - \bar{\rho}(r)). \quad (8)$$

Thus the two equations (7) and (8) form a system of differential equations in $P(r)$ and $\bar{\rho}(r)$.

For any physically reasonable astrophysical object, the pressure and density must be monotonically decreasing functions of the object's radius. Hence negativity of $P'(r)$ from (7) implies

$$\Lambda < 4\pi \bar{\rho}_b, \quad (9)$$

since by virtue of (6) the mean density is a monotonically decreasing function having its minimum at the boundary of the stellar object, where it is just the mean density of the stellar object. The boundary is defined by the radius R for which the pressure vanishes, $P(R) = 0$.

If, for the moment, we assume $\rho(r) = \rho = \text{const.}$, then (7) gives

$$P'(r) = -r \frac{\rho}{3} (4\pi \rho - \Lambda), \quad (10)$$

and is, as before, well defined if and only if $\Lambda < 4\pi\rho$. However, if $\Lambda = 4\pi\rho$ we find that the pressure must be constant for all r . Hence one may say that this is the Newtonian analogue of the Einstein static universe. Note that the cosmological constant in the Newtonian case is independent of the pressure as is expected from general grounds since in Newton's theory of gravity pressure does not contribute to the energy momentum tensor.

Moreover if the pressure increases near the center, it automatically diverges. This changes considerably in the general relativistic case.

3 General relativistic hydrostatic equilibrium

Most parts of the former section can still be applied to the general relativistic case. The only difference is that the differential equation for the pressure (7) is

replaced by the Tolman-Oppenheimer-Volkoff equation [38,39] with cosmological constant (TOV- Λ) and is given as

$$P'(r) = -r \frac{\rho(r)}{3} \left(1 + \frac{\lambda P(r)}{\rho(r)} \right) \left(\frac{\lambda 12\pi P(r) + 4\pi \bar{\rho}(r) - \Lambda}{1 - \lambda \frac{8\pi}{3} \bar{\rho}(r) r^2 - \lambda \frac{\Lambda}{3} r^2} \right), \quad (11)$$

where the velocity of light has been included by $\lambda = 1/c^2$ to facilitate the comparison with the Newtonian case. The Newtonian limit [40], $\lambda \rightarrow 0$, of (11) reproduces (7). We will use geometrical units henceforth.

The TOV- Λ equation together with (8) form a system of differential equations if an equation of state $\rho = \rho(P)$ is specified.

The right hand side of (11) is well defined if the denominator is

$$y^2 := 1 - \frac{8\pi}{3} \bar{\rho}(r) r^2 - \frac{\Lambda}{3} r^2 \geq 0 \quad \forall r, \quad (12)$$

where we introduced a new variable y . The interesting question is whether one can find a general relativistic analogue of the upper bound of the cosmological constant (9). The answer to this question is affirmative, and it is related to fact that the TOV- Λ equation be well defined.

One can derive [27, 42, 43] the following inequality

$$y \geq \frac{12\pi P(r) + 4\pi \bar{\rho}(r) - \Lambda}{12\pi P_c + 4\pi \bar{\rho}_c - \Lambda}, \quad (13)$$

which can be used to arrive at the desired equation. Since the boundary of any astrophysical object is defined by $P = 0$, the variable y is well defined up to the boundary if the remaining terms of the numerator satisfy

$$\Lambda < 4\pi \bar{\rho}_b. \quad (14)$$

Therefore, the Newtonian upper bound of the cosmological constant could exactly be reproduced in the general relativistic case.

Before continuing, let us shortly review qualitatively how one derives (13). The Einstein field equations for a static and spherically symmetric perfect fluid are three independent differential equations for four free functions. Hence one can specify one of them. The most physical one is to prescribe an equation of state $\rho = \rho(P)$. The TOV- Λ equation can be extracted from this system by eliminating the free function g_{rr} if the line element is given in its usual form.

¹ On the other hand, one can also eliminate the pressure from the second field equation and the conservation equation. After introducing Buchdahl variables [42] and assuming that the pressure is a decreasing function near the center and prescribing a monotonic equation of state one arrives at (13). Without the cosmological constant one can directly read it off from [42,43]. The cosmological constant is incorporated by using the standard substitution $\rho \rightarrow \rho + \Lambda/8\pi$ and

¹ $ds^2 = -e^{\nu(r)} dt^2 + e^{\alpha(r)} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2)$.

$P \rightarrow P - \Lambda/8\pi$ which leads again to equation (13). For an explicit derivation with the cosmological constant see [27, p.42–p.47].

One must be slightly careful with the above, since in case of a constant density distribution $\bar{\rho} = \rho_0$ one can integrate the TOV- Λ equation exactly and no upper bound $\Lambda = 4\pi\bar{\rho}$ arises, which is expected since the Einstein static universe with non-vanishing pressure is also a solution of the field equations. The interior metric has the geometry of a three sphere which is conformally flat [44] and one just has to deal with a harmless coordinate singularity. See [28] for a complete discussion of constant density solutions with cosmological constant where it is also shown that in contrast to the Newtonian case one can construct solutions with increasing pressure near the center.

4 Buchdahl inequalities

There are essentially two ways of solving the field equations, as already mentioned in the previous section. With the help of Buchdahl variables [42] one has a very convenient scheme to compare a general monotonically decreasing solution with a constant density solution. The latter is defined by the boundary of the mean density of the general solution.

To perform the described procedure, Buchdahl [42] originally assumed the existence of a global solution, which was later proven to exist in [43]. Up to $\Lambda < 4\pi\bar{\rho}_b$ one can use the methods of [43] to prove the existence of a global solution with non-vanishing cosmological constant [27, Theorem 5].

The derivation of the Buchdahl inequality is quite involved, therefore we only state the result and refer to [42] and [45, p.367] without and [27, Theorem 6] with the cosmological constant. The generalized Buchdahl inequality is given by

$$\sqrt{1 - \frac{8\pi}{3}\bar{\rho}_b R^2 - \frac{\Lambda}{3}R^2} \geq \frac{1}{3} - \frac{\Lambda}{12\pi\bar{\rho}_b}. \quad (15)$$

For constant density solutions, generalized Buchdahl inequalities can be found in [28, 46]. In [46] the surface redshift with Λ was derived from (15).

For $\Lambda = 0$ this inequality reduces to

$$\sqrt{1 - \frac{8\pi}{3}\bar{\rho}_b R^2} \geq \frac{1}{3}, \quad (16)$$

which with $M = (4\pi/3)\bar{\rho}R^3$ leads to its most popular form

$$\sqrt{1 - \frac{2M}{R}} \geq \frac{1}{3}, \quad 2M < \frac{8}{9}R, \quad (17)$$

from which one reads off the well known result that radii of static perfect fluid spheres are larger than the Schwarzschild radius of the corresponding mass.

Let us for the moment assume that a global solution exists independent of the cosmological constant, so we assume existence for $\Lambda \geq 4\pi\bar{\rho}_b$. From the generalized Buchdahl inequality (15) one finds instead

$$R^2 \leq \frac{\frac{1}{3}\left(4 - \frac{\Lambda}{4\pi\bar{\rho}_b}\right)}{4\pi\bar{\rho}_b}, \quad (18)$$

which after some algebra can be written as

$$\frac{\Lambda}{16\pi\bar{\rho}_b} \leq 1 - \frac{9M}{4R}. \quad (19)$$

Since the right-hand side is bounded by 1, one simply finds

$$\Lambda \leq 16\pi\bar{\rho}_b, \quad (20)$$

which is in full agreement with the results from constant density solutions [28], where solutions exist up to a cosmological constant Λ_S , given by

$$\Lambda_S = 16\pi\bar{\rho} + 24P_c. \quad (21)$$

Therefore it seems that properties of the constant density solutions in [28] are also a feature of general solutions with any given equation of state. However, existence of solutions with $\Lambda \geq 4\pi\bar{\rho}_b$ so far could not be proved. Nonetheless *we conjecture their existence based on the generalized Buchdahl inequality*, similar to Buchdahl's original assumption. Moreover one may use $\bar{\rho}_b = 3M/(4\pi R^3)$ in (18) which yields

$$3M \leq \frac{2}{3}R + R\sqrt{\frac{4}{9} - \frac{\Lambda}{3}R^2}, \quad (22)$$

and reduces to (17) for $\Lambda = 0$. The above equation may also be regarded as the generalized Buchdahl inequality since it gives an upper bound on the mass. It is tempting to make a Taylor expansion with respect to Λ in equation (22), trying to extract some further information. However, by looking at (18) and (19) one sees that nothing can be found since both equations are linear in Λ . Moreover it should be noted that the square root appearing on the right hand side of (22) is well defined if

$$R \leq \sqrt{\frac{4}{3} \frac{1}{\sqrt{\Lambda}}}, \quad (23)$$

is satisfied by the object's radius R . Inserting the highest possible radius (23) in (22) we obtain

$$\frac{2}{3}\sqrt{\frac{4}{3} \frac{1}{\sqrt{\Lambda}}} \geq 3M. \quad (24)$$

It is now worthwhile to compare the inequalities (23) and (24) with constraints arising from the Newtonian limit. The Newtonian approximation has, of course, a certain range of validity. In the static, non-relativistic limit, the requirement of weak fields results into two strong inequalities of the form

$$\sqrt{\frac{6}{\Lambda}} \gg R \gg M, \quad M_{\max} = \frac{2}{3}\sqrt{2}\frac{1}{\sqrt{\Lambda}} \gg M, \quad (25)$$

valid up to small corrections of the order M/M_{\max} and $(M/M_{\max})^2$. What is striking, is the similarity of inequalities (23) and (24) to the restrictions (25) of the Newtonian limit. Disregarding different numerical factors which are of the order of unity and the fact that (25) are strong inequalities, we see that both are essentially the same. Inequalities (23) and (24) can be understood as constraints on Λ to keep the object in hydrostatic equilibrium from which they are derived. On the other hand, (25) ensures that the gravitational fields are not too strong. Hence in both cases one expects a restriction on a positive cosmological constant. What is surprising, however, is the fact that these restrictions are so similar in both cases. We will discuss further relevance of equations (23)-(25) in section 8.

5 Lessons from constant density solutions

In this section, static and spherically symmetric constant density perfect fluids are shortly revisited. A new inequality is derived which in the constant density case is shown to reduce to an equality. In the subsequent section this is used to extract information on the cosmological constant which is encoded in a cubic equation in Λ .

5.1 With equation of state

Assume than a constant density and an equation of state are prescribed, for instance, the equation of state of a polytrope

$$P = \kappa\rho^\gamma. \quad (26)$$

In this case we have $P'(r) = 0$, that is, we also have a constant pressure (we can take this as an approximation in the sense that $P'(r) \approx 0$ for a slowly varying density or pressure profile). Using this in the TOV- Λ equation (11) leads to

$$P'(r) = -\frac{1}{3}r^2(\rho + P) \left(\frac{4\pi\bar{\rho} + 12\pi P - \Lambda}{1 - \frac{8\pi}{3}\bar{\rho}r^2 - \frac{\Lambda}{3}r^2} \right) = 0, \quad (27)$$

and implies an equation for the constant density ρ :

$$4\pi\rho + 12\pi\kappa\rho^\gamma - \Lambda = 0. \quad (28)$$

Note that with vanishing cosmological constant $\Lambda = 0$, there would not be any acceptable solution for a positive energy density. Thus, we can say that in the presence of a cosmological constant, a system with constant density and constant pressure can be in hydrostatical equilibrium; the Einstein static universe is exactly such a system. In the case with $\Lambda = 0$, such a system could not be in hydrostatical equilibrium; sooner or later the system will collapse under its own gravity. From (28) we then can find $\rho = \rho(\kappa, \gamma, \Lambda)$. To illustrate the consequence of equation (28) let us assume that $\gamma = 1$. It then follows that an astrophysical object with $\rho \sim \rho_{\text{crit}}$ is stable. Such stable objects would have the lowest possible density among stable astrophysical objects and could be e.g. superclusters.

5.2 Without equation of state

Let us assume that no equation of state is specified and consider the inequality

$$y(r) \geq \zeta(r) \left(\frac{12\pi P(r) + 4\pi\bar{\rho}(r) - \Lambda}{12\pi P_c + 4\pi\rho_c - \Lambda} \right). \quad (29)$$

The functions $y(r)$ and $\zeta(P(r))$ are the Buchdahl variables [42], already mentioned above:

$$\begin{aligned} \zeta(P(r)) &= \exp \left(- \int_{P_c}^{P(r)} \frac{dP}{P + \rho(P)} \right), \\ y(r) &= \sqrt{1 - \frac{8\pi}{3}\bar{\rho}(r)r^2 - \frac{\Lambda}{3}r^2}. \end{aligned} \quad (30)$$

For a constant density distribution one finds

$$\zeta(r) = \frac{P_c + \rho}{P(r) + \rho}, \quad (31)$$

and equation (29) becomes an equality

$$\sqrt{1 - \frac{8\pi}{3}\bar{\rho}r^2 - \frac{\Lambda}{3}r^2} = \frac{P_c + \rho}{P(r) + \rho} \left(\frac{12\pi P(r) + 4\pi\bar{\rho}(r) - \Lambda}{12\pi P_c + 4\pi\rho_c - \Lambda} \right). \quad (32)$$

Solving for the pressure and using $\rho_c = \bar{\rho} = \rho$ we find

$$P(r) = \rho \left[\frac{(1 - \beta)(P_c + \rho[1 - y(r)]) - 3P_c y(r)}{3P_c(y(r) - 1) + \rho(y(r)[1 - \beta] - 3)} \right], \quad (33)$$

where $\beta = \Lambda/4\pi\rho$ and moreover

$$P_c = \rho(1 - \beta) \frac{y(R) - 1}{3y(R) - (1 - \beta)}. \quad (34)$$

Finiteness of the central pressure implies that the denominator must be well defined, hence greater than zero. In the simplified constant density case this implies

$$y(R) > \frac{1}{3}(1 - \beta), \quad (35)$$

which is just the Buchdahl inequality (15).

To justify why in the constant density case the inequality (29) becomes an equality, one can compare with the solution of the TOV- Λ equation for constant density [28]. One finds

$$P(r) = \rho \frac{(\beta - 1) + Cy(r)}{3 - Cy(r)}, \quad (36)$$

where C is a constant of integration which can be fixed with the condition $P(R) = 0$. This yields

$$C = \frac{1 - \beta}{y(R)}. \quad (37)$$

On the other hand, at the center $r = 0$ we have

$$P_c = \rho \frac{(\beta - 1)(y(R) - 1)}{3y(R) - (1 - \beta)}, \quad (38)$$

and therefore the constant of integration becomes

$$C = \frac{3P_c + (1 - \beta)\rho}{P_c + \rho}. \quad (39)$$

Solving for $y(r)$ from (36) yields

$$y(r) = \frac{1}{C} \frac{\rho(1 - \beta) + 3P(r)}{P(r) + \rho}. \quad (40)$$

If we replace the constant C from (39) we obtain equation (32) and hence (33). This justifies the conversion from an inequality to an equality in (29) for constant density. Thus it is trivially fulfilled for constant density solutions.

6 General solutions with equation of state

In view of the above results, we can assume a variable density in order to find a more restrictive condition on Λ . First of all, inequality (29) is valid for any density profile; on the other hand, since it must be fulfilled at any r in order that the pressure be a decreasing function of the radius, we can evaluate it at the boundary $r = R$, defined by the condition $P(R) = 0$, which leads to

$$y(R) \geq \zeta(P(R)) \left(\frac{4\pi\bar{\rho}_b - \Lambda}{12\pi P_c + 4\pi\rho_c - \Lambda} \right) \geq \left(\frac{4\pi\bar{\rho}_b - \Lambda}{12\pi P_c + 4\pi\rho_c - \Lambda} \right). \quad (41)$$

In this case we can safely assume an equation of state in the form of (26), so that the central pressure is only a function of the central density. The function $\zeta(P)$ can be integrated and becomes a function of the central density when evaluated at the boundary

$$\zeta(P(R)) = \left(\kappa^\gamma \rho_c^{\gamma(\gamma-1)} + 1 \right)^{\frac{\gamma}{\gamma-1}} = \zeta(\rho_c). \quad (42)$$

Equation (41) then becomes a cubic equation for Λ , namely,

$$f(\Lambda) = a\Lambda^3 + b\Lambda^2 + c\Lambda + d \geq 0, \quad (43)$$

where the coefficients are given explicitly as

$$\begin{aligned} a &= -\frac{1}{3}R^3, & b &= 1 + 8\pi P_c R^2 - \frac{8\pi}{3}R^2(\bar{\rho} - \rho_c) - \zeta^2, \\ c &= -24\pi P_c - 48\pi^2 P_c^2 R^2 + 64\pi^2 \bar{\rho} P_c R^2 + 8\pi \zeta^2 \bar{\rho} - 8\pi \rho_c - 32\pi^2 P_c \rho_c R^2 \\ &\quad + \frac{64}{3}\pi^2 R^2 \bar{\rho} \rho_c - \frac{16}{3}\pi^2 R^2 \rho_c^2, \\ d &= 144\pi^2 P_c^2 - 384\pi^3 P_c^2 \bar{\rho} R^2 - 16\pi^2 \zeta^2 \bar{\rho}^2 + 96\pi^2 P_c \rho_c - 256\pi^3 P_c \bar{\rho} \rho_c R^2 \\ &\quad + 16\pi^2 \rho_c^2 - \frac{128}{3}\pi^3 \bar{\rho} \rho_c^2 R^2, \end{aligned} \quad (44)$$

and where $\bar{\rho} = \bar{\rho}_b = \bar{\rho}(R)$ is the object's mean density. Thus we have the following parameters R , ρ_c , P_c , $\bar{\rho}_b$, which can be related as follows: the central pressure is connected with the central density through the equation of state so that $P_c = \kappa \rho_c^\gamma$. The radius of the configuration is related with the mean density at the surface as

$$R^3 = \frac{3M}{4\pi \bar{\rho}_b}, \quad (45)$$

while the mass (4) is just the volume integral of the density

$$M = \int_0^R 4\pi s^2 \rho(s) ds. \quad (46)$$

We need the density profile and the total mean density (together with an equation of state) to solve the cubic equation (43) for Λ .

It is interesting to further exploit the inequality (41). Although it is quite involved to extract information on the cosmological constant one can solve it for the central energy density. This yields

$$\rho_c \geq \rho_{c,\min} = \frac{\bar{\rho} - \frac{\Lambda}{4\pi}}{y(R)} - 3P_c + \frac{\Lambda}{4\pi}, \quad (47)$$

giving a lower bound on the central energy density. This is not surprising because we have upper bounds on the boundary mean density. Since the energy

density for astrophysical models is a decreasing function of the radius, one must find some lower bound, that in particular extends the boundary mean density.

A clever choice of new variables could be of help in order to get information on the cosmological constant from (41). For example one could use the effective quantities mentioned in the derivation of equation (13). However, the numerator of (47) suggests a definition of type $\Delta = \bar{\rho} - \Lambda/4\pi$. In future we plan to explore the astrophysical significance of inequality (41) and other versions thereof.

7 Inequalities from virial theorem

In what follows, the non-relativistic virial theorem is recalled and an upper bound on the cosmological constant is derived with surprising similarities to those derived before.

Note that there exist several relativistic formulations of the virial theorem in literature [47–51], however, they are not complete generalizations since some restrictive assumptions are always needed in their formulation.

The standard non-relativistic virial theorem reads

$$\frac{d^2 I_{jk}}{dt^2} = 4K_{jk} + 2W_{jk} + \dots, \quad (48)$$

where I_{jk} is the inertial tensor, K_{jk} the kinetic and W_{jk} the gravitational potential tensor. The dots in (48) indicate other possible terms due to magnetic field, pressure etc. If an external force is exerted on the object, we have to add to the right hand side of equation (48) the term [52, p.280]

$$V_{jk} = -\frac{1}{2} \int \rho \left(x_k \frac{\partial \Phi_{\text{ext}}}{\partial x_j} + x_j \frac{\partial \Phi_{\text{ext}}}{\partial x_k} \right) d^3x, \quad (49)$$

where Φ_{ext} is the external potential and the case of a cosmological constant corresponds to

$$\Phi_{\text{ext}} = -\frac{1}{6} \Lambda r^2. \quad (50)$$

Therefore the new virial theorem which accounts for the cosmological constant [53] takes the form

$$\frac{d^2 I_{jk}}{dt^2} = 4K_{jk} + 2W_{jk} + \frac{2}{3} I_{jk} \Lambda + \dots. \quad (51)$$

This is, in principle, a differential equation for I_{jk} provided K_{jk} and W_{jk} are given or corresponding differential equations in these variables are derived. It is very often more convenient to consider a less demanding task by simply noting that the trace W of W_{jk} is negative whereas the trace K of K_{jk} is positive definite. Then the gravitational equilibrium i.e. $d^2 I_{jk}/dt^2 = 0$ yields the inequality

$$-\frac{1}{3} \Lambda I + |W| \geq 0, \quad (52)$$

where I denotes the trace of the inertial tensor I_{jk} . To appreciate the meaning of this inequality we specialize to the case of constant density $\bar{\rho} = \rho$. It is then easy to show that (52) takes the form

$$8\pi\rho \geq A\Lambda, \quad (53)$$

where the quantity A depends only on the geometry of the object under consideration. More specifically it reads

$$A = \frac{16\pi}{3} \frac{\int r^2 d^3x}{\int \frac{\Phi_N}{\rho} d^3x}, \quad (54)$$

where Φ_N is the Newtonian part of the non-relativistic gravitational potential. For spherically symmetric objects one easily calculates $A = 2$ and therefore the virial inequality is simply

$$4\pi\rho \geq \Lambda. \quad (55)$$

We already mentioned the similarity between Buchdahl's inequalities (which are a consequence of hydrostatic equilibrium) and the inequalities describing the validity of the Newtonian approximation. A second kind of such a coincidence (if one can call it coincidence) occurs also here with regard to equation (55), (9) and (14). To add to this coincidence we mention that the authors of [32] by demanding stability of circular orbits derived further the inequality

$$4\pi\rho \geq 4\pi\rho_c \geq \Lambda. \quad (56)$$

No doubt, all three inequalities originating from different premises have something in common. The similarity between the three results is not trivial. Certainly, the hydrostatic and gravitational equilibrium are intuitively related. However, the hydrostatic equilibrium necessarily involves pressure whereas (55) is derived without using this concept. Indeed, the virial equation for an object in gravitational equilibrium with pressure reads

$$\frac{1}{3}\Lambda I + W + 2K + 3PV + \dots = 0, \quad (57)$$

where the pressure P is constant over the volume V (note that P can have different signs depending whether the pressure is internal or external) and the dots indicate the presence of terms related to magnetic fields, rotational effects, etc. We think that this issue deserves a closer inspection.

With the help of the virial theorem we can show a drastic astrophysical effect of the cosmological constant. Inequalities like (52) and (53) are useful to check if a given astrophysical system is in gravitational equilibrium without the knowledge of the kinetic tensor or, which is the same, the average velocity $\langle v^2 \rangle$ of the components of the system. It is often of utmost interest to know this velocity. The knowledge of the shape of the object, its density profile and Λ allows one to calculate

$$\langle v^2 \rangle = \frac{|W|}{M} - \frac{8\pi}{3} \frac{\rho_{\text{vac}}}{M} I. \quad (58)$$

To appreciate the effect of Λ let us assume a constant density and the shape of the astrophysical object to be an ellipsoid. Then the W and I can be calculated analytically [52]. For the inertial integral I we get

$$I = \rho \frac{4}{15} a_1 a_2 a_3 (a_1^2 + a_2^2 + a_3^2), \quad (59)$$

where a_i are the axes of the ellipsoid. The mean velocity can be now written as

$$\langle v^2 \rangle_{\text{ellipsoid}} = \frac{32\pi}{45} \rho \rho_{\text{vac}} a_1 a_2 a_3 (a_1^2 + a_2^2 + a_3^2) \left(\frac{3\pi}{4} \Gamma_{\text{ellipsoid}} - 1 \right). \quad (60)$$

For the oblate case ($a_1 = a_2 > a_3$, $e = \sqrt{1 - a_3^2/a_1^2}$) one calculates

$$\Gamma_{\text{oblate}} = \frac{\left(\frac{a_3}{a_1}\right) \sin^{-1} e}{1 + \frac{1}{2} \left(\frac{a_3}{a_1}\right)^2 e}. \quad (61)$$

The prolate case ($a_1 = a_2 < a_3$, $e = \sqrt{1 - a_1^2/a_3^2}$) gives

$$\Gamma_{\text{prolate}} = \frac{\left(\frac{a_1}{a_3}\right)^3 \ln\left(\frac{1+e}{1-e}\right)}{1 + 2 \left(\frac{a_1}{a_3}\right)^2 e}. \quad (62)$$

The triaxial case ($a_1 > a_2 > a_3$, $\theta = \cos^{-1}(a_3/a_1)$, $k = \sqrt{(a_1^2 - a_2^2)/(a_1^2 - a_3^2)}$) is simply

$$\Gamma_{\text{triaxial}} = \frac{a_1 a_2 F(\theta, k)}{a_1^2 + a_2^2 + a_3^2}, \quad (63)$$

where $F(\theta, k)$ is the elliptic integral of the first kind. Note now that for a flattened prolate ellipsoid we can approximate

$$\Gamma_{\text{prolate}} \simeq \left(\frac{a_1}{a_3}\right)^3 \ln\left(\frac{1+e}{1-e}\right). \quad (64)$$

Since the nowadays preferred value of ρ_{vac} is $(0.7 - 0.8)\rho_{\text{crit}}$ we can say that if the constant ρ/ρ_{crit} is, say, 10^3 , it suffices for the ellipsoid to have the ratio $a_1/a_3 \sim 10$ in order that the mean velocity of its components approaches zero. This is valid always under the assumption that the object is in gravitational equilibrium. This effect is due to the relatively large cosmological constant. In general we can say that in flattened astrophysical systems in gravitational equilibrium, the mean velocity gets affected by the cosmological constant. The denser the system, the bigger should be the deviation from spherical symmetry to have a sizable effect. It is interesting if such an effect can be observed in reality which would confirm the existence of Λ . One can paraphrase this also by looking at the results from a different perspective. If we are certain that a given

astrophysical object is in gravitational equilibrium, then equation (52) and (53) would put a stringent bound on Λ in the case of strong deviation from spherical symmetry.

From the last equations we can infer that the power of equilibria concepts with Λ lies in analyzing relatively low density astrophysical objects whose shapes deviate from the spherical symmetry. It is evident that this can easily be done using the virial theorem. We have shown that for spherically symmetric objects the virial theorem and the hydrostatic equilibrium yield surprisingly similar results. It would be therefore desirable to have a tool to probe the hydrostatic equilibrium with Λ for arbitrarily shaped bodies as well. The full examination of this aspect of hydrostatic equilibrium goes beyond the scope of the present paper. We can, however, outline a direction in which such an investigation might proceed following [54]. The basic equations for the Newtonian concept of hydrostatic equilibrium are

$$\nabla P = -\rho \nabla \Phi, \quad \nabla^2 \Phi = 4\pi\rho - \Lambda, \quad (65)$$

together with (26) as the equation of state. Defining $u = \rho/\rho_0$, $\boldsymbol{\xi} = s\mathbf{x}$ with $s^2 = 4\pi\rho^{2-\gamma}/\kappa\gamma$ we obtain for $u = u(\boldsymbol{\xi})$

$$(1 - \gamma) \left(\frac{\nabla_{\boldsymbol{\xi}} u}{u} \right)^2 + \nabla_{\boldsymbol{\xi}}^2 \ln u = a_{\Lambda} u^{1-\gamma} - u^{2-\gamma}, \quad (66)$$

which is a generalized Lane-Emden equation. For $\Lambda = 0$ we have to set $a_{\Lambda} = 0$ whereas the case $\Lambda \neq 0$ requires $a_{\Lambda} = 2$. Furthermore for $\Lambda = 0$, ρ_0 is arbitrary, but is fixed to be $\rho_0 = \rho_{\text{vac}}$ for non-zero cosmological constant. We will examine the consequences of equation (66) elsewhere. However, here we can already mention a difference between the cases $\Lambda = 0$ and $\Lambda \neq 0$ due to the fact that the cosmological constant sets a scale for the density. Consider the case $\gamma = 1$. Then with $\gamma = 1$ and for $\Lambda = 0$ we have a scaling property in the following sense: if $u(\boldsymbol{\xi})$ is a solution of the equation, so is $\lambda^2 u(\lambda\boldsymbol{\xi})$ [55]. This similarity property is lost if $a_{\Lambda} = 2$.

Coming back to the virial theorem we note that interestingly one can eliminate the cosmological constant if we address the question whether an astrophysical object is in gravitational equilibrium. Let the system which is in gravitational equilibrium be described by I_1 , W_1 and K_1 , etc. This system serves us as a reference system. Defining $\epsilon_{21} \equiv I_2/I_1 = 1/\epsilon_{12}$ and provided $\Lambda \neq 0$, a second system (denoted by the subscript 2) is also in gravitational equilibrium if the following relation is fulfilled

$$\epsilon_{21} = \frac{W_2 + 2K_2 + \Xi_2}{W_1 + 2K_1 + \Xi_1}, \quad (67)$$

where Ξ_i contains terms related to pressure, magnetic field, etc. On the other hand, if we know two systems bounded by an external medium at a constant pressure P_{ext} (this means that $\Xi_i = -3P_{\text{ext}}V_i$ assuming that there is no magnetic

field) are in equilibrium, we can solve for the pressure and obtain

$$P_{\text{ext}} = \frac{1}{3V_1} \left(\frac{V_2}{V_1} - \epsilon_{21} \right)^{-1} \left[W_1 \left(\frac{W_2}{W_1} - \epsilon_{21} \right) + 2K_1 \left(\frac{K_2}{K_1} - \epsilon_{21} \right) \right]. \quad (68)$$

Note that the salient and necessary assumption for equations (67) and (68) is a non-zero cosmological constant Λ . With these equations we can put in relation two systems in equilibrium or solve for the variable common in both systems. Solving for the kinetic energy of system 2 we have

$$K_2 = \frac{1}{2} (2\epsilon_{21}K_1 + \epsilon_{21}W_1 - W_2 + \epsilon_{21}\Xi_1 - \Xi_2). \quad (69)$$

Since this is a positive definite quantity, we find an inequality for W_2

$$|W_2| \geq \epsilon_{21} \left(|W_1| - 2K_1 - \Xi_1 + \frac{1}{\epsilon_{21}}\Xi_2 \right), \quad (70)$$

which gives an estimate of the left-hand side.

As an example, we assume again that the system 1 is in steady state and that both systems are in equilibrium with an external pressure P_{ext} , as before. Under these conditions we can solve for the pressure and find

$$P_{\text{ext}} \leq \frac{|W_1|}{3V_1} \left(\frac{|W_2|}{|W_1|} - \epsilon_{21} \right) \left(\epsilon_{21} - \frac{V_2}{V_1} \right)^{-1}, \quad (71)$$

which can serve as a lower bound for the external pressure. The question that rises here is, what can we identify as the reference system? On the one hand, one could argue that at every different astrophysical scale there must be a standard system, so that one must find, for instance, a standard galaxy in order to obtain information about a certain other galaxy. The exploration of these interesting aspects will be the subject of further research. We conclude this section by the remark that the virial theorem with non-zero Λ has also been successfully applied in cosmology [56].

8 Scales of the cosmological constant

To appreciate the orders of magnitude estimate we reinstall here the Newtonian coupling constant G which we put to 1 in the preceding sections. The cosmological constant sets scales for density, length, mass and time. The density scale in equation (1) indicates the so-called coincidence problem. The latter can be formulated as the question as to why we are living in an epoch in which $\rho_{\text{vac}} \sim \rho_{\text{crit}}$. More precisely, since ρ_{crit} is epoch dependent, the coincidence $\rho_{\text{vac}} \sim \rho_{\text{crit}}$ is a time coincidence of a sharp transition from small ratio Ω_Λ to $\Omega_\Lambda \sim 1$ [57]. We can reformulate this coincidence in terms of length, noting that

$$r_\Lambda = \frac{1}{\sqrt{\Lambda}} = 0.476 H_0^{-1} = 1.43 \times 10^3 h_0^{-1} \text{Mpc}, \quad (72)$$

which is of cosmological order of magnitude and close to the radius of the visible universe. The combination

$$M_\Lambda = \frac{1}{G\sqrt{\Lambda}} = 3.61 \times 10^{22} h_0^{-1} M_\odot \sqrt{\frac{\rho_{\text{crit}}}{\rho_{\text{vac}}}}, \quad (73)$$

is close to the mass of the universe. As noted in [31] this is not trivial since Λ as a constant could be either smaller or bigger than what is assumed today. We could equally well live in an epoch where the universe is smaller or bigger or the mass of the universe is not close to M_Λ . Interestingly, both such scales play an important role in the considerations of the validity of the weak field approximation and in the hydrostatic equilibrium in the form of Buchdahl inequalities (see the discussion in section 4). Hereby the differences of the numerical factors are of order one. Had we lived in a different universe or a different epoch of our universe, the above scales would appear only in the Newtonian limit and the Buchdahl inequalities, which by itself is a remarkable technical coincidence. However, in our universe these scales appear in all considerations: cosmology, hydrostatic equilibrium and Newtonian limit. Notice also that one of the equations of motion of a test body in the Schwarzschild metric with Λ is the relation between the affine parameter λ and the time t which reads

$$\left(1 - 2\frac{r_S}{r} - \frac{1}{3}\frac{r^2}{r_\Lambda^2}\right) \frac{dt}{d\lambda} = E = \text{const.}, \quad (74)$$

with $r_S = GM$. The case $E = 0$ sets the limit of validity of the coordinate system. This means that the expression on the left hand side of (74) is zero which in turn leads to the solutions $r_{\text{min}} = 2r_S$ and $r_{\text{max}} = \sqrt{3}r_\Lambda$. Albeit the cases $r = r_{\text{min}}, r_{\text{max}}$ are singular points due to the choice of coordinates, the condition above tells us also that we should not go beyond the horizon of a black hole and also not beyond the horizon of the universe. The latter emerges only because of the above coincidence of length mentioned above. In the same context, i.e., considering the motion in a Schwarzschild spacetime the scale R defined by

$$R^3 = 3r_S r_\Lambda^2, \quad (75)$$

not only yields astrophysically relevant length scales (this is due to the combination of a small and large quantity), but also has the meaning of the largest possible radius within which bound orbits in the Schwarzschild metric are possible [58].

Curiously, not all scales are of cosmological or astrophysical relevance even though they have the right order of magnitude. The time scale connected to Λ is

$$T_\Lambda = 4.65 h_0^{-1} \text{Gyr}, \quad (76)$$

which is close to the age of our solar system, but should be considered as an accidental coincidence.

The scale $m_\Lambda = \sqrt{\Lambda} = 3.1 \times 10^{-42}$ GeV has no connection to particle physics, but interestingly establishes one more of Dirac's large numbers in the form $m_p/(G\sqrt{\Lambda}) \sim 3.2 \times 10^{41}$ where m_p is the nucleon's mass.

9 Summary

In this work we investigated the astrophysical limits on the cosmological constant for spherically symmetric bodies. These limits emerged from examinations of hydrostatic and gravitational equilibrium as well as from constraints on the validity of the Newtonian limit. Although the resulting inequalities are based on quite different premises, they do resemble each other to an extent which let us conjecture a deeper underlying principle or connection. As a side result we conjecture the existence of a new class of static and spherically symmetric perfect fluid solutions with the cosmological constant. By coincidence the length and mass scales set by Λ appear as the horizon and mass of the universe and simultaneously as the limits of validity of the weak field expansion and the hydrostatic equilibrium. This adds to the puzzle of the coincidence problem.

In this context we note that the inequalities based on the hydrostatic equilibrium were derived in the general relativistic framework whereas the virial inequalities required only a Newtonian approximation. We suspect that general relativistic virial equations might be a more suitable tool to compare the hydrostatic equilibrium with the virial one.

Already the Newtonian virial theorem is a powerful equation in many situations where the astrophysical object is not spherically symmetric. Indeed, flattened, large and diluted objects cannot be in gravitational equilibrium if the cosmological constant is as large as claimed nowadays. It is therefore of some interest to extend the consideration of the hydrostatic equilibrium allowing for an arbitrary shape of the objects. Unfortunately, the general relativistic treatment of arbitrarily shaped objects is quite out of reach at the moment, but we have indicated how to do that in the Newtonian approach. This will be covered elsewhere. Similarly, some of our results, like the inequality (43) still await its practical exploitation.

The inequality $2\rho \geq \rho_{\text{vac}}$ which we found from various considerations is valid provided that the object is spherically shaped and in equilibrium. Assuming this and taking into account that ρ can be at the most of the order of the background density i.e. ρ_{crit} , this is not a weak inequality. Indeed, it comes very close to the preferred $\rho_{\text{vac}} \sim 0.7\rho_{\text{crit}}$.

In this work we have laid down the basics of equilibria concepts including a cosmological constant. As indicated in the paper one should continue this study by examining hydrostatic equilibrium for non-spherical objects and generalize the equilibria concepts also to other theories of the Dark Energy [59].

We can look upon many inequalities derived in the main text by imposing the condition of equilibrium from two different angles. Assuming a known value of Λ the inequalities limit the possible values of the density of the astrophysical object if the latter is assumed in equilibrium. If we know that an object is

in equilibrium and assuming we can infer the mean density of such an object from some other considerations, then the same inequalities put a limit on the cosmological constant.

Acknowledgements

One of the authors CGB wishes to thank Herbert Balasin, Daniel Grumiller and Wolfgang Kummer for their constant support. This work has partly been supported by project P-15449-N08 of the Austrian Science Foundation (FWF) and is part of the research project Nr. 01/04 *Quantum Gravity, Cosmology and Categorification* of the Austrian Academy of Sciences (ÖAW) and the National Academy of Sciences of Ukraine (NASU).

References

- [1] L. M. Krauss. The End of the Age Problem, and the Case for a Cosmological Constant Revisited. *Astrophys. J.*, 501:461, 1998.
- [2] S. Perlmutter, M. S. Turner, and M. J. White. Constraining dark energy with sne ia and large-scale structure. *Phys. Rev. Lett.*, 83:670–673, 1999.
- [3] B. Ratra and P. J. E. Peebles. Cosmological consequences of a rolling homogeneous scalar field. *Phys. Rev.*, D37:3406, 1988.
- [4] C. Wetterich. Cosmology and the fate of dilatation symmetry. *Nucl. Phys.*, B302:668, 1988.
- [5] A. Yu. Kamenshchik, U. Moschella, and V. Pasquier. An alternative to quintessence. *Phys. Lett.*, B511:265–268, 2001.
- [6] G. M. Kremer and F. P. Devecchi. Viscous cosmological models and accelerated universes. *Phys. Rev.*, D67:047301, 2003.
- [7] G. M. Kremer. Irreversible processes in a universe modelled as a mixture of a chaplygin gas and radiation. *Gen. Rel. Grav.*, 35:1459–1466, 2003.
- [8] G. M. Kremer. Cosmological models described by a mixture of van der waals fluid and dark energy. *Phys. Rev.*, D68:123507, 2003.
- [9] G. M. Kremer and D. S. M. Alves. Acceleration field of a universe modeled as a mixture of scalar and matter fields. *Gen. Rel. Grav.*, 36:2039–2051, 2004.
- [10] S. Nojiri and S. D. Odintsov. Modified gravity with negative and positive powers of curvature: Unification of inflation and cosmic acceleration. *Phys. Rev.*, D68:123512, 2003.

- [11] Y. B. Zel'dovich. Cosmological constant and elementary particles. *JETP Lett.*, 6:316, 1967.
- [12] S. Carneiro. Holography and the cosmic coincidence. 2002. *gr-qc/0206064*.
- [13] M. Tegmark, M. Zaldarriaga, and A. J. S. Hamilton. Towards a refined cosmic concordance model: joint 11- parameter constraints from cmb and large-scale structure. *Phys. Rev.*, D63:043007, 2001.
- [14] R. A. J. Matthews. Dirac's coincidences 60 years on. *Astronomy and Geophysics*, 39:19–20, 1988.
- [15] A. Eddington. On the Value of the Cosmical Constant. *Royal Society of London Proceedings Series A*, 133:605–615, October 1931.
- [16] Mena M., Guillermo A., and S. Carneiro. Holography and the large number hypothesis. *Phys. Rev.*, D65:087303, 2002.
- [17] L. M. Krauss. The state of the universe: Cosmological parameters 2002. In *ESO-CERN Symposium on Astronomy, Cosmology and Fundamental Physics*, March 2003.
- [18] S. Perlmutter et al. Discovery of a supernova explosion at half the age of the universe and its cosmological implications. *Nature*, 391:51–54, 1998.
- [19] A. G. Riess et al. Type ia supernova discoveries at $z > 1$ from the hubble space telescope: Evidence for past deceleration and constraints on dark energy evolution. *Astrophys. J.*, 607:665–687, 2004.
- [20] S. W. Allen, R. W. Schmidt, H. Ebeling, A. C. Fabian, and L. van Speybroeck. Constraints on dark energy from chandra observations of the largest relaxed galaxy clusters. 2004. *astro-ph/0405340*.
- [21] M. Kunz and B. A. Bassett. A tale of two distances. 2004.
- [22] G. Gamov and E. Teller. On the Origin of Great Nebulae. *Phys. Rev.*, 55:654–657, 1939.
- [23] J. D. North. *The Measure of the Universe. A study of Modern Cosmology*. Dover Publications, 1990.
- [24] J. M. Maldacena. The large N limit of superconformal field theories and supergravity. *Adv. Theor. Math. Phys.*, 2:231–252, 1998.
- [25] M. S. Volkov, N. Straumann, George V. Lavrelashvili, M. Heusler, and O. Brodbeck. Cosmological analogues of the Bartnik–McKinnon solutions. *Phys. Rev.*, D54:7243–7251, 1996.
- [26] Y. Brihaye and B. Hartmann. Spherically symmetric solutions of a $(4+n)$ -dimensional Einstein-Yang-Mills model with cosmological constant. 2004.

- [27] C. G. Böhrer. General relativistic static fluid solutions with cosmological constant. 2002. *gr-qc/0308057*, *unpublished Diploma Thesis*.
- [28] C. G. Böhrer. Eleven spherically symmetric constant density solutions with cosmological constant. *Gen. Rel. Grav.*, 36:1039–1054, 2004.
- [29] Z. Stuchlík. Spherically symmetric static configurations of uniform density in spacetimes with a non-zero cosmological constant. *Acta Physica Slovaca*, 50:219–228, 2000.
- [30] H. Weyl. Über die statischen kugelsymmetrischen Lösungen von Einsteins «kosmologischen» Graviationsgleichungen. *Physikalische Zeitschrift*, 20:31–34, 1919.
- [31] M. Nowakowski. The consistent Newtonian limit of Einstein’s gravity with a cosmological constant. *Int. J. Mod. Phys.*, D10:649–662, 2001.
- [32] R. A. Sussman and X. Hernandez. On the Newtonian limit and cut-off scales of isothermal dark matter halos with cosmological constant. *Mon. Not. Roy. Astron. Soc.*, 345:871, 2003.
- [33] M. Nowakowski, J.-C. Sanabria, and A. Garcia. Gravitational equilibrium in the presence of a positive cosmological constant. *Phys. Rev.*, D66:023003, 2002.
- [34] C. G. Böhrer. The Einstein static universe with torsion and the sign problem of the cosmological constant. *Class. Quant. Grav.*, 21:1119–1124, 2004.
- [35] A. Dimakis and F. Mueller-Hoissen. Solutions of the Einstein-Cartan-Dirac equations with vanishing energy momentum tensor. *J. Math. Phys.*, 26:1040, 1985.
- [36] P. van Nieuwenhuizen. Supergravity. *Phys. Rept.*, 68:189–398, 1981.
- [37] S. Weinberg. *Gravitation and cosmology*. John Wiley & Sons, 1972.
- [38] J. R. Oppenheimer and G. M. Volkoff. On massive neutron cores. *Phys. Rev.*, 55:374–381, 1939.
- [39] R. C. Tolman. Static solutions of Einstein’s field equations for spheres of fluid. *Phys. Rev.*, 55:364–373, 1939.
- [40] J. Ehlers. *The Newtonian limit of general relativity*, page 112. In Ferrarese [41], 1991.
- [41] G. Ferrarese, editor. *Classical Mechanics and Relativity: Relationship and Consistency*. American Institute of Physics, 1991.
- [42] H. A. Buchdahl. General relativistic fluid spheres. *Phys. Rev.*, 116:1027–1034, 1959.

- [43] A. D. Rendall and B. G. Schmidt. Existence and properties of spherically symmetric static fluid bodies with a given equation of state. *Class. Quant. Grav.*, 8:985–1000, 1991.
- [44] H. Stephani. Über Lösungen der Einsteinschen Feldgleichungen, die sich in einen fünfdimensionalen flachen Raum einbetten lassen. *Commun. Math. Phys.*, 4:137–142, 1967.
- [45] R. Beig and B. G. Schmidt. Time-independent gravitational fields. *Lect. Notes Phys.*, 540:325–372, 2000.
- [46] M. K. Mak, Jr. Dobson, P. N., and T. Harko. Maximum mass-radius ratio for compact general relativistic objects in Schwarzschild-de Sitter geometry. *Mod. Phys. Lett.*, A15:2153–2158, 2000.
- [47] S. Bonazzola. The Virial Theorem in General Relativity. *Astrophys. J.*, 182:335–342, 1973.
- [48] S. Bonazzola and E.ourgoulhon. A virial identity applied to relativistic stellar models. *Class. Quant. Grav.*, 11:1775–1784, 1994.
- [49] A. Georgiou. A virial theorem for general relativistic charged fluids. *J. Phys.*, A13:3751–3759, 1980.
- [50] A. Georgiou. A virial theorem for rotating charged perfect fluids in general relativity. *Class. Quant. Grav.*, 20:359–367, 2003.
- [51] C. Vilain. Virial theorem in general relativity - Consequences for stability of spherical symmetry. *Astrophys. J.*, 227:307–318, 1979.
- [52] J. Binney and S. Tremaine. *Galactic Dynamics*. Princeton University Press, 1987. (Princeton Series in Astrophysics).
- [53] J. C. Jackson. The dynamics of clusters of galaxies in universes with non-zero cosmological constant, and the virial theorem mass discrepancy. *Mon. Not. Roy. Astron. Soc.* 148:249–260, 1970.
- [54] M. Lombardi and G. Bertin. Boyle’s law and gravitational instability. *Astronomy & Astrophysics*, 375:1091, 2001.
- [55] S. Chandrasekhar. *An Introduction to the Study of Stellar Structure*. Dover, 1967.
- [56] J. D. Barrow and G. Gotz. Newtonian no-hair theorems. *Class. Quant. Grav.*, 6:1253–1265, 1989.
- [57] P. H. Frampton. Dark energy: A pedagogic review. 2004.
- [58] A. Balaguera-Antolínez, C. G. Böhrer, and M. Nowakowski. Sales set by the cosmological constant. 2005. In preparation.
- [59] D. F. Mota and C. van de Bruck. On the spherical collapse model in dark energy cosmologies. *Astronomy & Astrophysics*, 421:71, 2004.